

Asymptotics of a Class of Fredholm Determinants

Craig A. Tracy

*Department of Mathematics and Institute of Theoretical Dynamics
University of California, Davis, CA 95616, USA
e-mail address: tracy@itd.ucdavis.edu*

Harold Widom

*Department of Mathematics
University of California, Santa Cruz, CA 95064, USA
e-mail address: widom@math.ucsc.edu*

Abstract

In this expository article we describe the asymptotics of certain Fredholm determinants which provide solutions to the cylindrical Toda equations, and we explain how these asymptotics are derived. The connection with Fredholm determinants arising in the theory of random matrices, and their asymptotics, are also discussed.

1. Introduction

In the Gaussian unitary ensemble, scaled in the bulk so that the density of eigenvalues becomes one, the probability $E(0; s)$ that an interval of length s contains no eigenvalues is equal to the Fredholm determinant of the operator on $L^2(0, s)$ with kernel

$$\frac{1}{\pi} \frac{\sin \pi(x - y)}{x - y}. \quad (1)$$

This is a result of M. Gaudin [6]. This quantity is directly related to the spacings between consecutive eigenvalues. The conditional probability density that, given that there is an eigenvalue at x , the next larger eigenvalue lies in an infinitesimal neighborhood of $x + s$ is equal to $-d^2 E(0; s)/ds^2$. This quantity is independent of x since the kernel depends only on the difference of the arguments.

To see how large the gaps between successive eigenvalues might be the asymptotics of $E(0; s)$ as $s \rightarrow \infty$ become of interest. (The asymptotics as $s \rightarrow 0$ are trivial.) A complete asymptotic expansion was found by F. J. Dyson [5], and reads

$$E(0; s/\pi) \sim 2^{1/3} e^{3\zeta'(-1)} s^{-1/4} e^{-s^2/8} \left(1 + \frac{a_1}{s} + \frac{a_2}{s^2} + \cdots \right),$$

with computable constants $a_1, a_2, \dots$. The main part of the asymptotics (all but the parenthetical correction terms) were obtained by applying a formal scaling argument to a result of the second author [17] on the asymptotics of certain Toeplitz determinants. The asymptotics were completed using inverse scattering techniques.

The derivation described above was not rigorous. The first rigorous result of this type is also due to the second author [18, 19] and was that $E(0; s) = e^{-s^2/8+o(s)}$. More exactly,

$$\frac{d}{ds} \log E(0; s) = -\frac{s}{4} + o(1).$$

This was subsequently extended to a complete asymptotic expansion by Deift, Its and Zhou [4]. Thus all of the Dyson expansion has now been proved, except for the constant factor.

Incidentally this constant is expressible in terms of the Barnes G -function [1], which will also arise below. This is an entire function satisfying $G(1) = 1$ and the functional equation

$$G(z+1) = \Gamma(z) G(z).$$

It turns out that

$$G\left(\frac{1}{2}\right)^2 = 2^{1/12} \pi^{-1/2} e^{3\zeta'(-1)}.$$

The kernel (1) arising in the Gaussian unitary ensemble has the form

$$\frac{\varphi(x) \psi(y) - \psi(x) \varphi(y)}{x - y}, \quad (2)$$

with $\varphi = \sin \pi x / \sqrt{\pi}$ and $\psi(x) = \cos \pi x \sqrt{\pi}$. Many other kernels of this same form arise in random matrix theory and for some there are (heuristically derived) asymptotic expansions analogous to the Dyson expansion [12, 13]. One of the interesting things about the function $E(0; s)$ is that it is expressible in terms of a Painlevé transcendent, more precisely P_V [8]. It was shown by the authors [12, 13, 14] that analogous quantities arising in several other matrix ensembles are also expressible in terms of Painlevé functions. All the kernels had the same general form (2), and this fact was crucial for the argument.

The first connection between Fredholm determinants and a Painlevé function was established in a paper of B. M. McCoy, T. T. Wu and the first author [10]. The function is the solution to the cylindrical sinh-Gordon equation (reducible to a special case of the P_{III} equation)

$$q''(t) + \frac{1}{t} q'(t) = 8 \sinh 2q \quad (3)$$

which satisfies

$$q(t) \sim \frac{\lambda}{\sqrt{2\pi t}} e^{-4t} \quad \text{as } t \rightarrow \infty.$$

In the cited work q was shown to be expressible as a certain sum of multiple integrals. This sum is in fact equal to a combination of logarithms of Fredholm determinants of two integral operators. Soon thereafter [11] an integral representation was found, in terms of this same function q , for $\det(I - \lambda^2 K^2)$, where K is the integral operator on $L^2(0, \infty)$ with kernel

$$\frac{e^{-t(x+x^{-1}+y+y^{-1})}}{x+y}. \quad (4)$$

This is also of the form (2). (More precisely it can be brought to that form with the change of variable $x \rightarrow x^{1/2}$.) More recently [3, 15] it was discovered that there is a direct

connection between this function and Fredholm determinants associated with K . This is $q = \log \det(I + \lambda K) - \log \det(I - \lambda K)$. (With hindsight it can be seen that this representation is equivalent to the one found in [10].) It is a generalization of the kernel (4) which we shall be considering here.

The *connection problem* for equations such as (3) is to determine the behavior of a singular point (in this case $t = 0$) if the behavior at another singular point (in this case $t = \infty$) is known. This connection problem was solved in [10]; the asymptotics will be a special case of those we shall describe below. The Dyson expansion, of course, solves the connection problem for another Painlevé function.

We shall describe here the results and methods of [16] which give the asymptotics of the Fredholm determinants of a class of kernels containing the above K as a special case. These determinants provide solutions to the cylindrical Toda equations

$$q_k''(t) + \frac{1}{t} q_k'(t) = 4(e^{q_k - q_{k-1}} - e^{q_{k+1} - q_k}), \quad k \in \mathbf{Z}. \quad (5)$$

If we have a solution q of (3) then $q_k = (-1)^k q$ is a solution to (5) which is 2-periodic in the sense that $q_{k+2} = q_k$ for all k . Conversely any 2-periodic solution of (5) gives the solutions $q = (q_k - q_{k-1})/2$ of (3).

We introduce a class of n -periodic solutions to the Toda equations generalizing the solution stated above for the sinh-Gordon equation. Define a family of kernels indexed by $k \in \mathbf{Z}$ by the formulas

$$K_k(x, y) = \sum_{\substack{\omega^n = 1 \\ \omega \neq 1}} c_\omega \omega^k \frac{e^{-t[(1-\omega)x + (1-\omega^{-1})x^{-1}]}}{-\omega x + y}.$$

This is clearly n -periodic in k , and it was shown in [20] that

$$q_k = \log \det(I - K_k) - \log \det(I - K_{k-1}) \quad (6)$$

provides a solution to (5). In the special case $n = 2$ there is a single coefficient and we recover the solution to (3) described above. (The exponential factors in the kernels are different, but the determinants are easily shown to be the same.)

It is the asymptotics as $t \rightarrow 0$ of these determinants $\det(I - K_k)$ which we shall discuss, and of course this will give the asymptotics of the solution q_k . The asymptotics take the form

$$\det(I - K_k) \sim b t^a, \quad (t \rightarrow 0),$$

where the constants a and b are determined by k and the coefficients c_ω . (We give the formulas for them and state our precise assumptions below.)

Asymptotic formulas of this type in the $n = 2$ case for $\det(I - K^2)$ were obtained earlier. The constant a was determined in [7] and b in [2]. In fact these were obtained for a more general class of kernels, where K^2 is replaced by KK^t and K now denotes the kernel (4) multiplied by a factor $(x/y)^\theta$. As was already mentioned, the asymptotics of q for the $n = 2$ case were found in [10]; in [9] a method was described to find connection formulas for solutions of a class of equations including some of the $n = 3$ cases of (5).

2. Statement of the results

Since K_k is obtained from K_0 by multiplying the coefficients c_ω by ω^k we may as well assume that $k = 0$. We do this and write K instead of K_0 . It is also convenient to change the constant b so that the asymptotic formula reads

$$\det(I - K) \sim b \left(\frac{t}{n}\right)^a, \quad (t \rightarrow 0).$$

The constants a and b are expressed in terms of certain zeros $\alpha_1, \dots, \alpha_n$ of the function

$$\sin \pi s - \pi \sum c_\omega (-\omega)^{s-1}.$$

(The exponential is determined by taking $|\arg(-\omega)| < \pi$.) The formulas are

$$a = \frac{1}{n} \sum_{i=1}^n \alpha_i^2 - \frac{(n+1)(2n+1)}{6},$$

$$b = \frac{\prod_{|j| < n} G\left(\frac{j}{n} + 1\right)^{n-|j|}}{\prod_{i,j=1}^n G\left(\frac{\alpha_i - \alpha_j}{n} + 1\right)}.$$

Here G is the Barnes G -function mentioned earlier.

To explain which zeros the α_i are, and the assumptions under which this has been proved, we introduce a parameter λ and the function

$$\sin \pi s - \lambda \pi \sum c_\omega (-\omega)^{s-1}. \quad (7)$$

Denote the zeros of this function by $\alpha_i(\lambda)$ ($i \in \mathbf{Z}$), where $\alpha_i(0) = i$. Then we take $\alpha_i = \alpha_i(1)$ for $i = 1, \dots, n$. Observe that the $\alpha_i(\lambda)$ are analytic functions of λ away from those values for which the function has multiple zeros, and we choose a path in $\mathbf{C}$ from $\lambda = 0$ to $\lambda = 1$ which avoids all such values. Our basic assumption is that there is a path in $\mathbf{C}$ from $\lambda = 0$ to $\lambda = 1$ such that everywhere on the path

$$\Re \alpha_i(\lambda) < \Re \alpha_{i+1}(\lambda), \quad |\Re \alpha_i(\lambda) - i| < 1.$$

(We believe that the second condition is redundant.)

From these formulas we can compute the asymptotics of our solutions (6) of (5). They are given by

$$q_k(t) = A \log \left(\frac{t}{n}\right) + \log B + o(1),$$

where for $k = 1, \dots, n$ the constants A and B are given by

$$A = 2(\alpha_k - k), \quad B = \prod_{1 \leq j < k} \frac{\Gamma\left(\frac{\alpha_j - \alpha_k}{n} + 1\right)}{\Gamma\left(\frac{\alpha_k - \alpha_j}{n}\right)} \prod_{k < j \leq n} \frac{\Gamma\left(\frac{\alpha_j - \alpha_k}{n}\right)}{\Gamma\left(\frac{\alpha_k - \alpha_j}{n} + 1\right)},$$

and for other values of k are given by periodicity.

In the special case $n = 2$ this gives the asymptotics for $\det(I \pm \lambda K)$, with K given by (4), and for the corresponding solutions of (3) for all λ except real λ satisfying $|\lambda| \geq 1$. Consequently in this case at least the range of validity is the right one since the asymptotics are quite different for these λ [10].

3. Outline of the derivation

If R_λ denotes the resolvent kernel of K , the kernel of $K(I - \lambda K)^{-1}$, then

$$\frac{d}{d\lambda} \log \det(I - \lambda K) = -\text{tr } R_\lambda.$$

Hence

$$\log \det(I - K) = - \int_0^1 \text{tr } R_\lambda d\lambda = - \int_0^1 d\lambda \int_0^\infty R_\lambda(x, x) dx,$$

where the path of integration can be taken to be that in our basic assumption. Now our kernel is given by

$$K(x, y) = \sum c_\omega \frac{e^{-t[(1-\omega)x + (1-\omega^{-1})x^{-1}]}}{-\omega x + y},$$

and we are looking for the asymptotics as $t \rightarrow 0$. Observe that as $t \rightarrow 0$,

$$e^{-t(1-\omega^{-1})x^{-1}} \rightarrow 1 \text{ uniformly for } x \geq 1, \quad e^{-t(1-\omega)x} \rightarrow 1 \text{ uniformly for } x \leq 1.$$

Therefore if we define kernels $K^\pm(x, y)$, still acting on $L^2(0, \infty)$, by

$$K^+(x, y) = \sum c_\omega \frac{e^{-t(1-\omega)x}}{-\omega x + y}, \quad K^-(x, y) = \sum c_\omega \frac{e^{-t(1-\omega^{-1})x^{-1}}}{-\omega x + y},$$

with corresponding resolvent kernels $R_\lambda^\pm(x, y)$ then we might hope that $R_\lambda(x, x)$ is close to $R_\lambda^+(x, x)$ when $x \geq 1$ and close to $R_\lambda^-(x, x)$ when $x \leq 1$. This is true, and using operator techniques it can be shown that

$$\int_0^\infty R_\lambda(x, x) dx = \int_1^\infty R_\lambda^+(x, x) dx + \int_0^1 R_\lambda^-(x, x) dx + o(1)$$

as $t \rightarrow 0$, uniformly in λ . This is heart of the argument. Once we have this approximation then everything is plain sailing, since the operators $K^\pm$ can be transformed into Wiener-Hopf operators, and so there are exact integral representations for their resolvent kernels.

To describe this transformation on K^+ , say, we observe that we may take $t = 1$ since the variable change $x \rightarrow x/t$ replaces K^+ by the same kernel with $t = 1$. Assuming this to be the case we make the further assumption that each $\Re \omega < 0$. The reason we can make this assumption is that in the present discussion the ω may be arbitrary complex numbers except for nonnegative real numbers (when the kernels cease to represent trace class operators) and that in the end everything will depend analytically on these ω . So that the formula for the resolvent derived with the restriction holds generally.

Let us define

$$A(x, u) = \sum c_\omega e^{-(1-\omega)x} e^{\omega x u}, \quad B(u, x) = e^{-xu}.$$

These are both kernels of bounded operators from $L^2(0, \infty)$ to $L^2(0, \infty)$ (here is where we use $\Re \omega < 0$) and

$$K^+(x, y) = \int_0^\infty A(x, u) B(u, y) du,$$

and so $K^+ = AB$. Hence (our assumption guarantees that $I - \lambda K^+$ is invertible)

$$R_\lambda^+ = AB(I - \lambda AB)^{-1} = A(I - \lambda BA)^{-1}B.$$

Therefore R_λ is known once $(I - \lambda BA)^{-1}$ is. But the operator BA on $L^2(0, \infty)$ has kernel

$$BA(u, v) = \sum \frac{c_\omega}{(u+1) - \omega(v+1)},$$

and under the change of variable $u \rightarrow e^\xi - 1$, $v \rightarrow e^\eta - 1$ the operator is transformed into the operator, also on $L^2(0, \infty)$, with kernel

$$\sum \frac{c_\omega}{e^{(\xi-\eta)/2} - \omega e^{-(\xi-\eta)/2}}.$$

This is a Wiener-Hopf operator (its kernel is a function of the difference of the arguments) and so its resolvent is expressible in terms of a factorization of its symbol, 1 minus λ times the Fourier transform of the function

$$\sum \frac{c_\omega}{e^{\xi/2} - \omega e^{-\xi/2}}.$$

After a change of variable the Fourier transform becomes a Mellin transform and the symbol becomes

$$1 - \frac{\lambda\pi}{\sin \pi s} \sum c_\omega (-\omega)^{s-1}.$$

This is how the function (7) arises and why everything is expressed in terms of it.

The symbol is factored into products and quotients of Γ -functions whose arguments involve the zeros $\alpha_i(\lambda)$. To see how the Barnes function appears consider an integral of the form

$$\int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} \frac{\Gamma(\frac{s}{n} + \beta)}{\Gamma(\frac{s}{n} + \beta)} f(s) ds, \quad (8)$$

where f has period n . (Integrals like this which involve the α_i arise in the computation of the constant b .) To evaluate this we integrate

$$\frac{G'(\frac{s}{n} + \beta)}{G(\frac{s}{n} + \beta)} f(s)$$

over the infinite rectangle with vertices $\frac{1}{2} \pm i\infty$ and $n + \frac{1}{2} \pm i\infty$. This is equal to (8) by the periodicity of f and the fact that

$$\frac{G'(\frac{s+n}{n} + \beta)}{G(\frac{s+n}{n} + \beta)} - \frac{G'(\frac{s}{n} + \beta)}{G(\frac{s}{n} + \beta)} = \frac{\Gamma'(\frac{s}{n} + \beta)}{\Gamma(\frac{s}{n} + \beta)}.$$

On the other hand this is also equal to

$$2\pi i \sum_{\alpha} \frac{G'(\frac{\alpha}{n} + \beta)}{G(\frac{\alpha}{n} + \beta)} \text{res}(f, \alpha),$$

where α runs over the poles of f inside the rectangle. In the integrals that arise these poles are exactly the points $\alpha_1, \dots, \alpha_n$, and so (8) is equal to

$$2\pi i \sum_{i=1}^n \frac{G'(\frac{\alpha_i}{n} + \beta)}{G(\frac{\alpha_i}{n} + \beta)} \operatorname{res}(f, \alpha_i(\lambda)).$$

Acknowledgements

The work of the first author was supported in part by the National Science Foundation through Grant DMS-9303413, and the work of the second author was supported in part by the National Science Foundation through Grant DMS-9424292.

References

- [1] E. W. Barnes, *The theory of the G-function*, Quart. J. Pure and Appl. Math. **31** (1900) 264–314.
- [2] E. L. Basor and C. A. Tracy, *Asymptotics of a tau function and Toeplitz determinants with singular generating functions*, Int. J. Mod. Phys. **A7** (1992) 83–107.
- [3] D. Bernard and A. LeClair, *Differential equations for sine-Gordon correlation functions at the free fermion point*. Nucl. Phys. **B426** [FS] (1994) 534–558.
- [4] P. Deift, A. Its and X. Zhou, *A Riemann-Hilbert approach to asymptotic problems arising in the theory of random matrix models, and also in the theory of integrable statistical mechanics*, Ann. of Math. **146** (1997) 149–235.
- [5] F. J. Dyson, *Fredholm determinants and inverse scattering problems*, Commun. Math. Phys. **47** (1976) 171–183.
- [6] M. Gaudin, *Sur la loi limite de l’espacement des valeurs propres d’une matrice aléatoire*, Nucl. Phys. **25** (1961) 447–458. [Reprinted in C. E. Porter, *Statistical Theory of Spectra: Fluctuations*, New York, Academic Press, 1965.]
- [7] M. Jimbo, *Monodromy problem and the boundary condition for some Painlevé equations*, Publ. RIMS, Kyoto Univ. **18** (1982) 1137–1161.
- [8] M. Jimbo, T. Miwa, Y. Môri and M. Sato, *Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent*, Physica **1D** (1980) 80–158.
- [9] A. V. Kitaev, *Method of isometric deformation for “degenerate” third Painlevé equation*, J. Soviet Math. **46** (1989) 2077–2082.
- [10] B. M. McCoy, C. A. Tracy and T. T. Wu, *Painlevé functions of the third kind*, J. Math. Phys. **18** (1977) 1058–1092.

- [11] M. Sato, T. Miwa and M. Jimbo, *Holonomic quantum fields, III, IV*, Publ. RIMS, Kyoto Univ. **15** (1979) 577–629, 871–972.
- [12] C. A. Tracy and H. Widom, *Level-spacing distributions and the Airy kernel*, Commun. Math. Phys. **159** (1994) 151–174.
- [13] C. A. Tracy and H. Widom, *Level-spacing distributions and the Bessel kernel*, Commun. Math. Phys. **161** (1994) 289–309.
- [14] C. A. Tracy and H. Widom, *Fredholm determinants, differential equations and matrix models*, Commun. Math. Phys. **163** (1994) 38–72.
- [15] C. A. Tracy and H. Widom, *Fredholm determinants and the mKdV/sinh-Gordon hierarchies*, Commun. Math. Phys. **179** (1996) 1–10.
- [16] C. A. Tracy and H. Widom, *Asymptotics of a class of solutions to the cylindrical Toda equations*, Commun. Math. Phys. to appear.
- [17] H. Widom, *The strong Szegő limit theorem for circular arcs*, Ind. U. Math. J. **21** (1971) 277–283.
- [18] H. Widom, *The asymptotics of a continuous analogue of orthogonal polynomials*, J. Approx. Th. **76** (1994) 51–64.
- [19] H. Widom, *Asymptotics for the Fredholm determinant of the sine kernel on a union of intervals*, Commun. Math. Phys. **171** (1995) 159–180.
- [20] H. Widom, *Some classes of solutions to the Toda lattice hierarchy*, Commun. Math. Phys. **184** (1997) 653–667.